

Spectral asymptotics on negatively curved
surfaces and hyperbolic dynamics

Iosif Polterovich

(Université de Montréal)

References:

H. Lapointe, I. P., Y. Safarov, *Average growth of the spectral function on a Riemannian manifold*, Comm. Partial Differential Equations, 34 (2009), no. 6, 581-615.

D. Jakobson, I. P., J. Toth, *A lower bound for the remainder in Weyl's law on negatively curved surfaces*, Int. Math. Res. Not. (2008), no. 2, Art. ID rnm142, 38 pp.

D. Jakobson, I. P., *Estimates from below for the spectral function and for the remainder in local Weyl's law*, Geom. Funct. Anal. 17, no. 3 (2007), 806-838.

Preliminaries

Preliminaries

X^n , $n \geq 2$ — compact Riemannian manifold

Preliminaries

$X^n, n \geq 2$ — compact Riemannian manifold

Δ — Laplace operator

Preliminaries

X^n , $n \geq 2$ — compact Riemannian manifold

Δ — Laplace operator

$\Delta\phi_i = \lambda_i^2\phi_i$, $\{\phi_i\}$ — orthonormal basis of eigenfunctions

$0 < \lambda_1^2 \leq \lambda_2^2 \leq \dots$ — eigenvalues

Preliminaries

X^n , $n \geq 2$ — compact Riemannian manifold

Δ — Laplace operator

$\Delta\phi_i = \lambda_i^2\phi_i$, $\{\phi_i\}$ — orthonormal basis of eigenfunctions

$0 < \lambda_1^2 \leq \lambda_2^2 \leq \dots$ — eigenvalues

Spectral function:

$$N_{x,y}(\lambda) = \sum_{\lambda_i < \lambda} \phi_i(x)\phi_i(y)$$

Preliminaries

X^n , $n \geq 2$ — compact Riemannian manifold

Δ — Laplace operator

$\Delta\phi_i = \lambda_i^2\phi_i$, $\{\phi_i\}$ — orthonormal basis of eigenfunctions

$0 < \lambda_1^2 \leq \lambda_2^2 \leq \dots$ — eigenvalues

Spectral function:

$$N_{x,y}(\lambda) = \sum_{\lambda_i < \lambda} \phi_i(x)\phi_i(y)$$

Counting function:

$$N(\lambda) = \#\{\lambda_i < \lambda\} = \int_M N_{x,x}(\lambda)$$

Weyl's law

Weyl's law

Pointwise Weyl's law:

$$N_{x,x}(\lambda) = \frac{\lambda^n}{(4\pi)^{\frac{n}{2}} \Gamma(\frac{n}{2} + 1)} + R_x(\lambda),$$

where $R_x(\lambda) = O(\lambda^{n-1})$.

Weyl's law

Pointwise Weyl's law:

$$N_{x,x}(\lambda) = \frac{\lambda^n}{(4\pi)^{\frac{n}{2}} \Gamma(\frac{n}{2} + 1)} + R_x(\lambda),$$

where $R_x(\lambda) = O(\lambda^{n-1})$.

Integrating over the manifold X we get:

Weyl's law

Pointwise Weyl's law:

$$N_{x,x}(\lambda) = \frac{\lambda^n}{(4\pi)^{\frac{n}{2}} \Gamma(\frac{n}{2} + 1)} + R_x(\lambda),$$

where $R_x(\lambda) = O(\lambda^{n-1})$.

Integrating over the manifold X we get:

$$N(\lambda) = \frac{\text{Vol}(X) \lambda^n}{(4\pi)^{\frac{n}{2}} \Gamma(\frac{n}{2} + 1)} + R(\lambda),$$

where $R(\lambda) = O(\lambda^{n-1})$.

Weyl's law

Pointwise Weyl's law:

$$N_{x,x}(\lambda) = \frac{\lambda^n}{(4\pi)^{\frac{n}{2}} \Gamma(\frac{n}{2} + 1)} + R_x(\lambda),$$

where $R_x(\lambda) = O(\lambda^{n-1})$.

Integrating over the manifold X we get:

$$N(\lambda) = \frac{\text{Vol}(X) \lambda^n}{(4\pi)^{\frac{n}{2}} \Gamma(\frac{n}{2} + 1)} + R(\lambda),$$

where $R(\lambda) = O(\lambda^{n-1})$.

Remainder estimates are **sharp** and attained on a round sphere.

Main results

Main results

Lower bounds for $N_{x,y}(\lambda)$, $R_x(\lambda)$ and $R(\lambda)$.

Main results

Lower bounds for $N_{x,y}(\lambda)$, $R_x(\lambda)$ and $R(\lambda)$.

Two types of estimates: for arbitrary manifolds and for negatively curved manifolds.

Main results

Lower bounds for $N_{x,y}(\lambda)$, $R_x(\lambda)$ and $R(\lambda)$.

Two types of estimates: for arbitrary manifolds and for negatively curved manifolds.

When the curvature is negative, hyperbolic dynamics allows to prove finer estimates.

General case: off-diagonal bound

General case: off-diagonal bound

It was proved by **Lapointe–P.–Safarov** '09 that (under some generic technical condition)

General case: off-diagonal bound

It was proved by **Lapointe–P.–Safarov** '09 that (under some generic technical condition)

$$\frac{1}{\lambda} \int_0^\lambda |N_{x,y}(\mu)| d\mu \gg \lambda^{\frac{n-1}{2}}, \quad x \neq y;$$

General case: off-diagonal bound

It was proved by **Lapointe–P.–Safarov** '09 that (under some generic technical condition)

$$\frac{1}{\lambda} \int_0^\lambda |N_{x,y}(\mu)| d\mu \gg \lambda^{\frac{n-1}{2}}, \quad x \neq y;$$

This bound is expected to be always true. It is sharp and attained on a round sphere.

General case: off-diagonal bound

It was proved by **Lapointe–P.–Safarov** '09 that (under some generic technical condition)

$$\frac{1}{\lambda} \int_0^\lambda |N_{x,y}(\mu)| d\mu \gg \lambda^{\frac{n-1}{2}}, \quad x \neq y;$$

This bound is expected to be always true. It is sharp and attained on a round sphere.

Moreover, it is shown in **L.–P.–S.** that the spectral function grows *on average* as $\lambda^{\frac{n-1}{2}}$ on *any* manifold.

General case: on-diagonal bounds

General case: on-diagonal bounds

Similarly, the following lower bound is typically (and possibly always) true on any [surface](#):

$$\frac{1}{\lambda} \int_0^\lambda |R_x(\mu)| d\mu \gg \sqrt{\lambda},$$

General case: on-diagonal bounds

Similarly, the following lower bound is typically (and possibly always) true on any [surface](#):

$$\frac{1}{\lambda} \int_0^\lambda |R_x(\mu)| d\mu \gg \sqrt{\lambda},$$

In dimensions $n > 2$, one typically has

$$\frac{1}{\lambda} \int_0^\lambda |R_x(\mu)| d\mu \gg \lambda^{n-2},$$

General case: on-diagonal bounds

Similarly, the following lower bound is typically (and possibly always) true on any [surface](#):

$$\frac{1}{\lambda} \int_0^\lambda |R_x(\mu)| d\mu \gg \sqrt{\lambda},$$

In dimensions $n > 2$, one typically has

$$\frac{1}{\lambda} \int_0^\lambda |R_x(\mu)| d\mu \gg \lambda^{n-2},$$

The latter bound is simpler and proved using different techniques than the estimate for surfaces.

General case: bound on the error term in Weyl's law

General case: bound on the error term in Weyl's law

Under some mild assumptions, one can show that the error term $R(\lambda)$ satisfies

$$\frac{1}{\lambda} \int_0^\lambda |R(\mu)| d\mu \gg \lambda^{n-2}.$$

General case: bound on the error term in Weyl's law

Under some mild assumptions, one can show that the error term $R(\lambda)$ satisfies

$$\frac{1}{\lambda} \int_0^\lambda |R(\mu)| d\mu \gg \lambda^{n-2}.$$

Open question: Is

$$\limsup_{\lambda \rightarrow \infty} |R(\lambda)| = \infty$$

on any surface?

General case: bound on the error term in Weyl's law

Under some mild assumptions, one can show that the error term $R(\lambda)$ satisfies

$$\frac{1}{\lambda} \int_0^\lambda |R(\mu)| d\mu \gg \lambda^{n-2}.$$

Open question: Is

$$\limsup_{\lambda \rightarrow \infty} |R(\lambda)| = \infty$$

on any surface?

We give a **positive** answer to this question in the **negatively curved** case.

Manifolds of negative curvature

Manifolds of negative curvature

Suppose sectional curvatures satisfy

$$-K_1^2 \leq K(\xi, \eta) \leq -K_2^2$$

Manifolds of negative curvature

Suppose sectional curvatures satisfy

$$-K_1^2 \leq K(\xi, \eta) \leq -K_2^2$$

Let G^t be the *geodesic flow* on the unit tangent bundle SX .

Manifolds of negative curvature

Suppose sectional curvatures satisfy

$$-K_1^2 \leq K(\xi, \eta) \leq -K_2^2$$

Let G^t be the *geodesic flow* on the unit tangent bundle SX .

It is an **Anosov flow**, i.e. there exists natural splitting of $T_\xi(SX)$ into a direct sum of DG^t -invariant subspaces

$$T_\xi(SX) = E_\xi^u \oplus E_\xi^o \oplus E_\xi^s.$$

Manifolds of negative curvature

Suppose sectional curvatures satisfy

$$-K_1^2 \leq K(\xi, \eta) \leq -K_2^2$$

Let G^t be the *geodesic flow* on the unit tangent bundle SX . It is an **Anosov flow**, i.e. there exists natural splitting of $T_\xi(SX)$ into a direct sum of DG^t -invariant subspaces

$$T_\xi(SX) = E_\xi^u \oplus E_\xi^o \oplus E_\xi^s.$$

Here E_ξ^u is the **unstable** (exponentially expanding) subspace of dimension $(n - 1)$, E_ξ^s is the **stable** (exponentially contracting) subspace of dimension $(n - 1)$, and E_ξ^o is a one-dimensional subspace **tangent** to the flow.

Thermodynamic formalism for Anosov flows

Thermodynamic formalism for Anosov flows

Sinai-Ruelle-Bowen potential: $\mathcal{H} : SX \rightarrow \mathbf{R} :$

Thermodynamic formalism for Anosov flows

Sinai-Ruelle-Bowen potential: $\mathcal{H} : SX \rightarrow \mathbf{R} :$

$$\mathcal{H}(\xi) = \left. \frac{d}{dt} \right|_{t=0} \ln \det dG^t|_{E_\xi^u}.$$

Thermodynamic formalism for Anosov flows

Sinai-Ruelle-Bowen potential: $\mathcal{H} : SX \rightarrow \mathbf{R} :$

$$\mathcal{H}(\xi) = \left. \frac{d}{dt} \right|_{t=0} \ln \det dG^t|_{E_\xi^u}.$$

Topological pressure of $f : SX \rightarrow \mathbf{R}$:

Thermodynamic formalism for Anosov flows

Sinai-Ruelle-Bowen potential: $\mathcal{H} : SX \rightarrow \mathbf{R}$:

$$\mathcal{H}(\xi) = \left. \frac{d}{dt} \right|_{t=0} \ln \det dG^t|_{E_\xi^u}.$$

Topological pressure of $f : SX \rightarrow \mathbf{R}$:

$$P(f) = \sup_{\mu} \left(h_{\mu} + \int f d\mu \right),$$

μ is a G^t -invariant measure, h_{μ} — **Kolmogorov–Sinai entropy**.

Thermodynamic formalism for Anosov flows

Sinai-Ruelle-Bowen potential: $\mathcal{H} : SX \rightarrow \mathbf{R}$:

$$\mathcal{H}(\xi) = \left. \frac{d}{dt} \right|_{t=0} \ln \det dG^t|_{E_\xi^u}.$$

Topological pressure of $f : SX \rightarrow \mathbf{R}$:

$$P(f) = \sup_{\mu} \left(h_{\mu} + \int f d\mu \right),$$

μ is a G^t -invariant measure, h_{μ} — **Kolmogorov–Sinai entropy**.

Example $P(-\mathcal{H}) = 0$.

Variational principle: $P(0) = h$, h — **topological entropy** of G^t .

Variational principle: $P(0) = h$, h — **topological entropy** of G^t .

Margulis's theorem:

$$\nu(T) = \frac{e^{hT}}{hT}(1 + o(1)),$$

$\nu(T)$ — number of closed geodesics on X of length $\leq T$.

Local results: off-diagonal

Local results: off-diagonal

Theorem A (Jakobson–P.) If X is negatively curved then for any $\delta > 0$ and $x \neq y$

$$N_{x,y}(\lambda) \neq o\left(\lambda^{\frac{n-1}{2}} (\log \lambda)^{\frac{P(-\mathcal{H}/2)}{h} - \delta}\right)$$

Local results: off-diagonal

Theorem A (Jakobson–P.) If X is negatively curved then for any $\delta > 0$ and $x \neq y$

$$N_{x,y}(\lambda) \neq o\left(\lambda^{\frac{n-1}{2}} (\log \lambda)^{\frac{P(-\mathcal{H}/2)}{h} - \delta}\right)$$

Power of the logarithm is **positive**

$$\frac{P(-\mathcal{H}/2)}{h} \geq \frac{K_2}{2K_1},$$

Local results: off-diagonal

Theorem A (Jakobson–P.) If X is negatively curved then for any $\delta > 0$ and $x \neq y$

$$N_{x,y}(\lambda) \neq o\left(\lambda^{\frac{n-1}{2}} (\log \lambda)^{\frac{P(-\mathcal{H}/2)}{h} - \delta}\right)$$

Power of the logarithm is **positive**

$$\frac{P(-\mathcal{H}/2)}{h} \geq \frac{K_2}{2K_1},$$

and equals $\frac{1}{2}$ if curvature is *constant*.

Local results: off-diagonal

Theorem A (Jakobson–P.) If X is negatively curved then for any $\delta > 0$ and $x \neq y$

$$N_{x,y}(\lambda) \neq o\left(\lambda^{\frac{n-1}{2}} (\log \lambda)^{\frac{P(-\mathcal{H}/2)}{h} - \delta}\right)$$

Power of the logarithm is **positive**

$$\frac{P(-\mathcal{H}/2)}{h} \geq \frac{K_2}{2K_1},$$

and equals $\frac{1}{2}$ if curvature is *constant*.

Get a **log-improvement** compared to the general case!

Local results: on-diagonal

Theorem B (Jakobson–P.) Let X be negatively curved and $n = 2, 3$.

Local results: on-diagonal

Theorem B (Jakobson–P.) Let X be negatively curved and $n = 2, 3$. Then for any $\delta > 0$ and for any $x \in X$,

$$R_x(\lambda) \neq o\left(\lambda^{\frac{n-1}{2}} (\log \lambda)^{\frac{P(-\mathcal{H}/2)}{h} - \delta}\right).$$

Local results: on-diagonal

Theorem B (Jakobson–P.) Let X be negatively curved and $n = 2, 3$. Then for any $\delta > 0$ and for any $x \in X$,

$$R_x(\lambda) \neq o\left(\lambda^{\frac{n-1}{2}} (\log \lambda)^{\frac{P(-\mathcal{H}/2)}{h} - \delta}\right).$$

For $n = 2$, related bounds were proved by **Karnaukh** '96.

Local results: on-diagonal

Theorem B (Jakobson–P.) Let X be negatively curved and $n = 2, 3$. Then for any $\delta > 0$ and for any $x \in X$,

$$R_x(\lambda) \neq o\left(\lambda^{\frac{n-1}{2}} (\log \lambda)^{\frac{P(-\mathcal{H}/2)}{h} - \delta}\right).$$

For $n = 2$, related bounds were proved by **Karnaukh** '96.

If $n \geq 4$, we do not get anything new compared to the general case:

$$\frac{1}{\lambda} \int_0^\lambda |R_x(\mu)| d\mu \gg \lambda^{n-2}.$$

Remainder estimate on surfaces

Theorem C (Jakobson–P.–Toth) Let X be a compact surface of negative curvature. Then for any $\delta > 0$

$$R(\lambda) \neq o\left((\log \lambda)^{\frac{P(-\mathcal{H}/2)}{h} - \delta}\right).$$

Remainder estimate on surfaces

Theorem C (Jakobson–P.–Toth) Let X be a compact surface of negative curvature. Then for any $\delta > 0$

$$R(\lambda) \neq o\left((\log \lambda)^{\frac{P(-\mathcal{K}/2)}{h} - \delta}\right).$$

For *constant* negative curvature such a bound was proved in **1976** by **Randol** and **Hejhal** using methods of *analytic number theory*.

Remainder estimate on surfaces

Theorem C (Jakobson–P.–Toth) Let X be a compact surface of negative curvature. Then for any $\delta > 0$

$$R(\lambda) \neq o\left((\log \lambda)^{\frac{P(-\mathcal{H}/2)}{h} - \delta}\right).$$

For *constant* negative curvature such a bound was proved in **1976** by **Randol** and **Hejhal** using methods of *analytic number theory*.

Our approach: wave trace asymptotics for long times, thermodynamic formalism, small-scale microlocalization.

Theorem C is in agreement with the following

Conjecture (*folklore*) On a *generic* negatively curved surface

$$R(\lambda) = O(\lambda^\varepsilon)$$

for any $\varepsilon > 0$.

Sketch of the proof of Theorem C

Sketch of the proof of Theorem C

Consider the spectral distribution

$$e(t) = \sum_{i=0}^{\infty} \cos(\lambda_i t)$$

which is the even part of the wave trace on X .

Sketch of the proof of Theorem C

Consider the spectral distribution

$$e(t) = \sum_{i=0}^{\infty} \cos(\lambda_i t)$$

which is the even part of the wave trace on X .

One can show that $\text{sing supp } e(t) = \{0\} \cup \{\pm\sigma_l\}$, where σ_l denotes the **length spectrum** of X .

Sketch of the proof of Theorem C

Consider the spectral distribution

$$e(t) = \sum_{i=0}^{\infty} \cos(\lambda_i t)$$

which is the even part of the wave trace on X .

One can show that $\text{sing supp } e(t) = \{0\} \cup \{\pm\sigma_l\}$, where σ_l denotes the **length spectrum** of X .

Example On the unit circle

$$\sum_{k=1}^{\infty} \cos k t = -\frac{1}{2} + \pi \sum_{k=-\infty}^{\infty} \delta(x - 2 k \pi)$$

To prove Theorem C we use **long time** version of the **Duistermaat–Guillemin** trace formula.

To prove Theorem C we use **long time** version of the **Duistermaat–Guillemin** trace formula.

Plan: counting function \longleftrightarrow Fourier transform of the wave trace

To prove Theorem C we use **long time** version of the **Duistermaat–Guillemin** trace formula.

Plan: counting function \longleftrightarrow Fourier transform of the wave trace
Main term \longleftrightarrow singularity at $t = 0$

To prove Theorem C we use **long time** version of the **Duistermaat–Guillemin** trace formula.

Plan: counting function \longleftrightarrow Fourier transform of the wave trace

Main term \longleftrightarrow singularity at $t = 0$

Error term \longleftrightarrow singularities at $t \in \sigma_I$.

To prove Theorem C we use **long time** version of the **Duistermaat–Guillemin** trace formula.

Plan: counting function \longleftrightarrow Fourier transform of the wave trace

Main term \longleftrightarrow singularity at $t = 0$

Error term \longleftrightarrow singularities at $t \in \sigma_I$.

Singularities \longleftrightarrow growth of the Fourier transform

To prove Theorem C we use **long time** version of the **Duistermaat–Guillemin** trace formula.

Plan: counting function \longleftrightarrow Fourier transform of the wave trace

Main term \longleftrightarrow singularity at $t = 0$

Error term \longleftrightarrow singularities at $t \in \sigma_I$.

Singularities \longleftrightarrow growth of the Fourier transform

Originally, Duistermaat–Guillemin formula takes into account only **one** closed geodesic.

To prove Theorem C we use **long time** version of the **Duistermaat–Guillemin** trace formula.

Plan: counting function \longleftrightarrow Fourier transform of the wave trace

Main term \longleftrightarrow singularity at $t = 0$

Error term \longleftrightarrow singularities at $t \in \sigma_I$.

Singularities \longleftrightarrow growth of the Fourier transform

Originally, Duistermaat–Guillemin formula takes into account only **one** closed geodesic.

Let us sum up the contributions of **all** closed geodesics of length $L \leq T(\lambda)$, where $T(\lambda) \rightarrow \infty$ as $\lambda \rightarrow \infty$.

Notation

Notation

$\chi(t, T)$ — a cut-off function

$$\chi(t, T) = (1 - \psi(t)) \hat{\rho}\left(\frac{t}{T}\right),$$

Notation

$\chi(t, T)$ — a cut-off function

$$\chi(t, T) = (1 - \psi(t)) \hat{\rho}\left(\frac{t}{T}\right),$$

ρ is an even, non-negative **Schwartz** function, $\text{supp } \hat{\rho} \subset [-1, +1]$,

Notation

$\chi(t, T)$ — a cut-off function

$$\chi(t, T) = (1 - \psi(t)) \hat{\rho}\left(\frac{t}{T}\right),$$

ρ is an even, non-negative **Schwartz** function, $\text{supp } \hat{\rho} \subset [-1, +1]$,

$\psi(t) \in C_0^\infty(\mathbf{R})$, $\psi(t) \equiv 1$ if $t \in [-T_0, T_0]$, $\psi(t) \equiv 0$ if $|t| \geq 2T_0$.

Notation

$\chi(t, T)$ — a cut-off function

$$\chi(t, T) = (1 - \psi(t)) \hat{\rho}\left(\frac{t}{T}\right),$$

ρ is an even, non-negative **Schwartz** function, $\text{supp } \hat{\rho} \subset [-1, +1]$,

$\psi(t) \in C_0^\infty(\mathbf{R})$, $\psi(t) \equiv 1$ if $t \in [-T_0, T_0]$, $\psi(t) \equiv 0$ if $|t| \geq 2T_0$.

L_γ — **length** of a closed geodesic γ

Notation

$\chi(t, T)$ — a cut-off function

$$\chi(t, T) = (1 - \psi(t)) \hat{\rho}\left(\frac{t}{T}\right),$$

ρ is an even, non-negative **Schwartz** function, $\text{supp } \hat{\rho} \subset [-1, +1]$,

$\psi(t) \in C_0^\infty(\mathbf{R})$, $\psi(t) \equiv 1$ if $t \in [-T_0, T_0]$, $\psi(t) \equiv 0$ if $|t| \geq 2T_0$.

L_γ — **length** of a closed geodesic γ

L_γ^\sharp — **primitive period** of γ

Notation

$\chi(t, T)$ — a cut-off function

$$\chi(t, T) = (1 - \psi(t)) \hat{\rho}\left(\frac{t}{T}\right),$$

ρ is an even, non-negative **Schwartz** function, $\text{supp } \hat{\rho} \subset [-1, +1]$,

$\psi(t) \in C_0^\infty(\mathbf{R})$, $\psi(t) \equiv 1$ if $t \in [-T_0, T_0]$, $\psi(t) \equiv 0$ if $|t| \geq 2T_0$.

L_γ — **length** of a closed geodesic γ

L_γ^\sharp — **primitive period** of γ

\mathcal{P}_γ — **linearized Poincaré map** corresponding to γ

Proposition 1

Let $T(\lambda) \rightarrow \infty$ as $\lambda \rightarrow \infty$ with $T(\lambda) \leq \varepsilon \log \lambda$ for some $\varepsilon > 0$ small enough. Then the smoothed Fourier transform of the wave trace

$$w(\lambda, T) = \int_{-\infty}^{\infty} \chi(t, T) e(t) \cos \lambda t dt$$

has the following asymptotics:

Proposition 1

Let $T(\lambda) \rightarrow \infty$ as $\lambda \rightarrow \infty$ with $T(\lambda) \leq \varepsilon \log \lambda$ for some $\varepsilon > 0$ small enough. Then the smoothed Fourier transform of the wave trace

$$w(\lambda, T) = \int_{-\infty}^{\infty} \chi(t, T) e(t) \cos \lambda t dt$$

has the following asymptotics:

$$w(\lambda, T) = \sum_{L_\gamma \leq T(\lambda)} \frac{L_\gamma^\sharp \cos(\lambda L_\gamma) \chi(L_\gamma, T)}{\sqrt{|\det(I - \mathcal{P}_\gamma)|}} + O(1).$$

We also have

Proposition 2 Assume $R(\lambda) = o((\log \lambda)^a)$ for some $a > 0$.

We also have

Proposition 2 Assume $R(\lambda) = o((\log \lambda)^a)$ for some $a > 0$. Then $w(\lambda, T) = o((\log \lambda)^a)$.

We also have

Proposition 2 Assume $R(\lambda) = o((\log \lambda)^a)$ for some $a > 0$. Then $w(\lambda, T) = o((\log \lambda)^a)$.

Using Proposition 1 we show that $w(\lambda, T)$ grows sufficiently fast, and then apply Proposition 2 to prove Theorem C **by contradiction**.

Proposition 1: idea of the proof

Proposition 1: idea of the proof

First try: L_γ 's are the **singularities** of the wave trace. Apply stationary phase at each singularity and sum up.

Proposition 1: idea of the proof

First try: L_γ 's are the **singularities** of the wave trace. Apply stationary phase at each singularity and sum up.

Problem: As $T(\lambda) \rightarrow \infty$, singularities may **accumulate**, and a priori there is no control at which rate: "*almost multiplicities*" in the length spectrum can be arbitrarily close! It is impossible to separate L_γ 's, and the stationary phase method fails.

Proposition 1: idea of the proof

First try: L_γ 's are the **singularities** of the wave trace. Apply stationary phase at each singularity and sum up.

Problem: As $T(\lambda) \rightarrow \infty$, singularities may **accumulate**, and a priori there is no control at which rate: "*almost multiplicities*" in the length spectrum can be arbitrarily close! It is impossible to separate L_γ 's, and the stationary phase method fails.

Instead: **Microlocalize** and separate closed geodesics in the **phase space**.

Proposition 1: idea of the proof

First try: L_γ 's are the **singularities** of the wave trace. Apply stationary phase at each singularity and sum up.

Problem: As $T(\lambda) \rightarrow \infty$, singularities may **accumulate**, and a priori there is no control at which rate: "*almost multiplicities*" in the length spectrum can be arbitrarily close! It is impossible to separate L_γ 's, and the stationary phase method fails.

Instead: Microlocalize and separate closed geodesics in the **phase space**. To do this, we use the following dynamical result.

“Spaghetti lemma”

“Spaghetti lemma”

Let X be a negatively curved surface and let $\Omega(\gamma, d)$ denote the d -neighborhood of a geodesic γ in SX .

“Spaghetti lemma”

Let X be a negatively curved surface and let $\Omega(\gamma, d)$ denote the d -neighborhood of a geodesic γ in SX .

There exist positive constants T_0, B and τ such that for any $T > T_0$ the sets $\Omega(\gamma, e^{-BT})$ are disjoint for all pairs of closed geodesics γ on X with length $L_\gamma \in [T - \tau, T]$.

“Spaghetti lemma”

Let X be a negatively curved surface and let $\Omega(\gamma, d)$ denote the d -neighborhood of a geodesic γ in SX .

There exist positive constants T_0, B and τ such that for any $T > T_0$ the sets $\Omega(\gamma, e^{-BT})$ are disjoint for all pairs of closed geodesics γ on X with length $L_\gamma \in [T - \tau, T]$.

Proof uses uniqueness of a closed geodesic in each free homotopy class.

Growth of $w(\lambda, T)$

By Proposition 1

$$w(\lambda, T) = \sum_{L_\gamma \leq T(\lambda)} \frac{L_\gamma^\sharp \cos(\lambda L_\gamma) \chi(L_\gamma, T)}{\sqrt{|\det(I - \mathcal{P}_\gamma)|}} + O(1).$$

Growth of $w(\lambda, T)$

By Proposition 1

$$w(\lambda, T) = \sum_{L_\gamma \leq T(\lambda)} \frac{L_\gamma^\# \cos(\lambda L_\gamma) \chi(L_\gamma, T)}{\sqrt{|\det(I - \mathcal{P}_\gamma)|}} + O(1).$$

Because of $\cos(\lambda L_\gamma)$, it is an **oscillating sum!** We need to estimate it from **below**.

First, let us forget about oscillations.

First, let us forget about oscillations.

Consider the sum

$$\sum_{L_\gamma \leq T} \frac{L_\gamma}{\sqrt{|\det(I - \mathcal{P}_\gamma)|}}$$

First, let us forget about oscillations.

Consider the sum

$$\sum_{L_\gamma \leq T} \frac{L_\gamma}{\sqrt{|\det(I - \mathcal{P}_\gamma)|}}$$

To find the asymptotics of this sum we use the following property of the **topological pressure** $P(f)$ due to **Parry** and **Pollicott**:

$$\sum_{L_\gamma \leq T} L_\gamma \exp \left[\int_\gamma f(\gamma(s), \gamma'(s)) ds \right] \sim \frac{C e^{P(f)T}}{P(f)}.$$

Proposition 3

There exists a constant $C_0 > 0$ such that

$$\sum_{L_\gamma \leq T} \frac{L_\gamma}{\sqrt{|\det(I - \mathcal{P}_\gamma)|}} = C_0 e^{P(-\frac{\mathcal{H}}{2}) \cdot T} (1 + o(1))$$

as $T \rightarrow \infty$, where P is the topological pressure and \mathcal{H} is the Sinai-Ruelle-Bowen potential.

Proposition 3

There exists a constant $C_0 > 0$ such that

$$\sum_{L_\gamma \leq T} \frac{L_\gamma}{\sqrt{|\det(I - \mathcal{P}_\gamma)|}} = C_0 e^{P(-\frac{\mathcal{H}}{2}) \cdot T} (1 + o(1))$$

as $T \rightarrow \infty$, where P is the topological pressure and \mathcal{H} is the Sinai-Ruelle-Bowen potential.

One can show that $P(-\frac{\mathcal{H}}{2}) \geq \frac{K_2}{2} > 0$ and hence the sum grows **exponentially** in T .

Proposition 3

There exists a constant $C_0 > 0$ such that

$$\sum_{L_\gamma \leq T} \frac{L_\gamma}{\sqrt{|\det(I - \mathcal{P}_\gamma)|}} = C_0 e^{P(-\frac{\mathcal{H}}{2}) \cdot T} (1 + o(1))$$

as $T \rightarrow \infty$, where P is the topological pressure and \mathcal{H} is the Sinai-Ruelle-Bowen potential.

One can show that $P(-\frac{\mathcal{H}}{2}) \geq \frac{K_2}{2} > 0$ and hence the sum grows **exponentially** in T .

Now we deal with the oscillations in $w(\lambda, T)$.

Dirichlet box principle

Dirichlet box principle

One can choose λ large enough so that

$$\text{dist}(\lambda L_\gamma, 2\pi\mathbf{Z}) < \frac{1}{10}$$

for **all** $L_\gamma \leq T$.

Dirichlet box principle

One can choose λ large enough so that

$$\text{dist}(\lambda L_\gamma, 2\pi\mathbf{Z}) < \frac{1}{10}$$

for **all** $L_\gamma \leq T$.

Then $\cos(\lambda L_\gamma)$ is always close to 1 for all L_γ .

Dirichlet box principle

One can choose λ large enough so that

$$\text{dist}(\lambda L_\gamma, 2\pi\mathbf{Z}) < \frac{1}{10}$$

for **all** $L_\gamma \leq T$.

Then $\cos(\lambda L_\gamma)$ is always close to 1 for all L_γ .

This “**straightening the phases**” argument is well-known in analytic number theory.

By Margulis's theorem,

$$\#\{\gamma \mid L_\gamma \leq T\} \sim \frac{e^{hT}}{hT},$$

where h is the topological entropy.

By Margulis's theorem,

$$\#\{\gamma \mid L_\gamma \leq T\} \sim \frac{e^{hT}}{hT},$$

where h is the topological entropy.

To make Dirichlet box principle work one has to take

$$T \approx \frac{1}{h} \log \log \lambda$$

By Margulis's theorem,

$$\#\{\gamma \mid L_\gamma \leq T\} \sim \frac{e^{hT}}{hT},$$

where h is the topological entropy.

To make Dirichlet box principle work one has to take

$$T \approx \frac{1}{h} \log \log \lambda$$

Thus, exponential growth in Proposition 3 yields a **logarithmic** lower bound in Theorem C. □