

La mesure de Mahler supérieure et la question de Lehmer

Matilde Lalín

(joint with Kaneenika Sinha (IISER, Kolkata))

Université de Montréal

mlalin@dms.umontreal.ca

<http://www.dms.umontreal.ca/~mlalin>

Conférence Québec-Maine

Université Laval

2 octobre, 2010

Mahler measure for one-variable polynomials

Pierce (1918): $P \in \mathbb{Z}[x]$ monic,

$$P(x) = \prod_i (x - r_i)$$

$$\Delta_n = \prod_i (r_i^n - 1)$$

$$P(x) = x - 2 \Rightarrow \Delta_n = 2^n - 1$$

Lehmer (1933):

$$\lim_{n \rightarrow \infty} \frac{|r^{n+1} - 1|}{|r^n - 1|} = \begin{cases} |r| & \text{if } |r| > 1 \\ 1 & \text{if } |r| < 1 \end{cases}$$

For

$$P(x) = a \prod_i (x - r_i)$$

$$M(P) = |a| \prod_{|r_i| > 1} |r_i|, \quad m(P) = \log |a| + \sum_{|r_i| > 1} \log |r_i|.$$

By Jensen's formula,

$$m(P) := \int_0^1 \log \left| P \left(e^{2\pi i \theta} \right) \right| d\theta.$$

Kronecker's Lemma

$P \in \mathbb{Z}[x]$, $P \neq 0$,

$$m(P) = 0 \Leftrightarrow P(x) = x^k \prod \Phi_{n_i}(x)$$

where Φ_{n_i} are cyclotomic polynomials.

Lehmer's question

Lehmer (1933):

Given $\epsilon > 0$, can we find a polynomial $P(x) \in \mathbb{Z}[x]$ such that $0 < m(P) < \epsilon$?

$$m(x^{10} + x^9 - x^7 - x^6 - x^5 - x^4 - x^3 + x + 1) \cong 0.162357612\dots$$

Is the above polynomial the best possible?

$$\sqrt{\Delta_{379}} = 1,794,327,140,357$$

Lehmer's question

Lehmer (1933):

Given $\epsilon > 0$, can we find a polynomial $P(x) \in \mathbb{Z}[x]$ such that $0 < m(P) < \epsilon$?

$$m(x^{10} + x^9 - x^7 - x^6 - x^5 - x^4 - x^3 + x + 1) \cong 0.162357612\dots$$

Is the above polynomial the best possible?

$$\sqrt{\Delta_{379}} = 1,794,327,140,357$$

$P \in \mathbb{C}[x]$ reciprocal iff

$$P(x) = \pm x^{\deg P} P(x^{-1}).$$

Theorem (Smyth, 1971)

$P \in \mathbb{Z}[x]$ nonreciprocal,

$$m(P) \geq m(x^3 - x - 1) \cong 0.2811995743 \dots$$

Higher Mahler measure

For $k \in \mathbb{Z}_{\geq 0}$, the k -higher Mahler measure of P is

$$m_k(P) := \int_0^1 \log^k \left| P(e^{2\pi i \theta}) \right| d\theta.$$

$$k = 1 : \quad m_1(P) = m(P),$$

and

$$m_0(P) = 1.$$

The simplest examples

(Kurokawa, L., Ochiai)

$$m_2(x-1) = \frac{\zeta(2)}{2} = \frac{\pi^2}{12}.$$

$$m_3(x-1) = -\frac{3\zeta(3)}{2}.$$

$$m_4(x-1) = \frac{3\zeta(2)^2 + 21\zeta(4)}{4} = \frac{19\pi^4}{240}.$$

$$m_5(x-1) = -\frac{15\zeta(2)\zeta(3) + 45\zeta(5)}{2}.$$

$$m_6(x-1) = \frac{45}{2}\zeta(3)^2 + \frac{275}{1344}\pi^6.$$

$$\begin{aligned}
\sum_{k=0}^{\infty} \frac{m_k(x-1)}{k!} s^k &= \int_0^1 |e^{2\pi i\theta} - 1|^s d\theta \\
&= \frac{\Gamma(s+1)}{\Gamma\left(\frac{s}{2}+1\right)^2} = \exp\left(\sum_{k=2}^{\infty} \frac{(-1)^k(1-2^{1-k})\zeta(k)}{k} s^k\right).
\end{aligned}$$

Transform the integral into a Beta function, then use the Weierstrass product of the Γ -function.

Lehmer's question for even higher measure

Theorem

If $P(x) \in \mathbb{Z}[x]$, then for any $h \geq 1$,

$$m_{2h}(P) \geq \begin{cases} \left(\frac{\pi^2}{12}\right)^h, & \text{if } P(x) \text{ is reciprocal,} \\ \left(\frac{\pi^2}{48}\right)^h, & \text{if } P(x) \text{ is non-reciprocal.} \end{cases}$$

- $m_2(P)$ for P reciprocal is minimized when P is a product of monomials and cyclotomic polynomials.
- $m_2(P) \geq \frac{\pi^2}{12}$ for P a product of monomials and cyclotomic polynomials.
- For P nonreciprocal, take $P(x)x^{\deg P}P(x^{-1})$.
- For $m_{2h}(P)$, use Hölder's inequality.

Multiple Mahler measure

Let $P_1, \dots, P_k \in \mathbb{C}[x^\pm]$ be non-zero Laurent polynomials. Their multiple higher Mahler measure is defined by

$$m(P_1, \dots, P_k) = \int_0^1 \log |P_1(e^{2\pi i\theta})| \cdots \log |P_k(e^{2\pi i\theta})| d\theta.$$

$$m_2 \left(\prod_i (x - r_i) \right) = \sum_{i,j} m(x - r_i, x - r_j).$$

Multiple Mahler measure for simple polynomials

Lemma (Kurokawa, L., Ochiai)

For $0 \leq \alpha \leq 1$

$$m(x - 1, x - e^{2\pi i\alpha}) = \frac{\pi^2}{2} \left(\alpha^2 - \alpha + \frac{1}{6} \right).$$

m_2 of cyclotomic polynomials

Proposition

- For any two positive integers a and b ,

$$m(x^a - 1, x^b - 1) = \frac{\pi^2}{12} \frac{(a, b)^2}{ab}.$$

- For a positive integer n , let $\phi_n(x)$ denote the n -th cyclotomic polynomial and φ Euler's function. Then

$$m_2(\phi_n(x)) = \frac{\pi^2}{12} \frac{\varphi(n) 2^{r(n)}}{n},$$

where $r(n)$ denotes the number of distinct prime divisors of n .

m_2 of noncyclotomic polynomials

| $P(x)$ | $m(P)$ | $m_2(P)$ |
|--|---------------------|---------------------|
| $x^8 + x^5 - x^4 + x^3 + 1$ | 0.2473585132 | 1.0980813745 |
| $x^{10} + x^9 - x^7 - x^6 - x^5 - x^4 - x^3 + x + 1$ | 0.1623576120 | 1.7447964556 |
| $x^{10} - x^6 + x^5 - x^4 + 1$ | 0.1958888214 | 1.2863292447 |
| $x^{10} + x^7 + x^5 + x^3 + 1$ | 0.2073323581 | 1.2320444893 |
| $x^{10} - x^8 + x^5 - x^2 + 1$ | 0.2320881973 | 1.1704950485 |
| $x^{10} + x^8 + x^7 + x^5 + x^3 + x^2 + 1$ | 0.2368364616 | 1.1914083866 |
| $x^{10} + x^9 - x^5 + x + 1$ | 0.2496548880 | 1.0309287773 |
| $x^{12} + x^{11} + x^{10} - x^8 - x^7 - x^6 - x^5 - x^4 + x^2 + x + 1$ | 0.2052121880 | 1.4738375004 |
| $x^{12} + x^{11} + x^{10} + x^9 - x^6 + x^3 + x^2 + x + 1$ | 0.2156970336 | 1.5143823478 |
| $x^{12} + x^{11} - x^7 - x^6 - x^5 + x + 1$ | 0.2239804947 | 1.2059443050 |
| $x^{12} + x^{10} + x^7 - x^6 + x^5 + x^2 + 1$ | 0.2345928411 | 1.2434560052 |
| $x^{12} + x^{10} + x^9 + x^8 + 2x^7 + x^6 + 2x^5 + x^4 + x^3 + x^2 + 1$ | 0.2412336268 | 1.6324129051 |
| $x^{14} + x^{11} - x^{10} - x^7 - x^4 + x^3 + 1$ | 0.1823436598 | 1.3885013172 |
| $x^{14} - x^{12} + x^7 - x^2 + 1$ | 0.1844998024 | 1.3845721865 |
| $x^{14} - x^{12} + x^{11} - x^9 + x^7 - x^5 + x^3 - x^2 + 1$ | 0.2272100851 | 1.4763006621 |
| $x^{14} + x^{11} + x^{10} + x^9 + x^8 + x^7 + x^6 + x^5 + x^4 + x^3 + 1$ | 0.2351686174 | 1.4352060397 |
| $x^{14} + x^{13} - x^8 - x^7 - x^6 + x + 1$ | 0.2368858459 | 1.2498299096 |
| $x^{14} + x^{13} + x^{12} - x^9 - x^8 - x^7 - x^6 - x^5 + x^2 + x + 1$ | 0.2453300143 | 1.3362661982 |
| $x^{14} + x^{13} - x^{11} - x^7 - x^3 + x + 1$ | 0.2469561884 | 1.3898540050 |

Known noncyclotomic $P \in \mathbb{Z}[x]$, $\deg P \leq 14$, $m(P) < 0.25$.

(Mossinghoff: <http://www.cecm.sfu.ca/~mjm/Lehmer/search/>)

Lemma (Kurokawa, L., Ochiai)

$$\begin{aligned} m(x-1, x-e^{2\pi i\alpha}, x-e^{2\pi i\beta}) &= -\frac{1}{4} \sum_{1 \leq k, l} \frac{\cos 2\pi((k+l)\beta - l\alpha)}{kl(k+l)} \\ &\quad -\frac{1}{4} \sum_{1 \leq k, m} \frac{\cos 2\pi((k+m)\alpha - m\beta)}{km(k+m)} \\ &\quad -\frac{1}{4} \sum_{1 \leq l, m} \frac{\cos 2\pi(l\alpha + m\beta)}{lm(l+m)}. \end{aligned}$$

$$m_3 \left(\frac{x^n - 1}{x - 1} \right) = \frac{3}{2} \zeta(3) \left(\frac{-2 + 3n - n^3}{n^2} \right) + \frac{3\pi}{2} \sum_{\substack{j=1 \\ n \nmid j}}^{\infty} \frac{\cot \left(\pi \frac{j}{n} \right)}{j^2}.$$

Examples

$$m_3 (x^2 + x + 1) = -\frac{10}{3} \zeta(3) + \frac{\sqrt{3}\pi}{2} L(2, \chi_{-3}).$$

$$m_3 (x^3 + x^2 + x + 1) = -\frac{81}{16} \zeta(3) + \frac{3\pi}{2} L(2, \chi_{-4}).$$

Lehmer's question for odd higher measure

Theorem

Let $P_n(x) = \frac{x^n - 1}{x - 1}$. For $h \geq 1$ fixed,

$$\lim_{n \rightarrow \infty} m_{2h+1}(P_n) = 0.$$

Moreover, the sequence $m_{2h+1}(P_n)$ is nonconstant.

Theorem (essentially Boyd, Lawton)

Let $P(x_1, \dots, x_n) \in \mathbb{C}[x_1, \dots, x_n]$ and $\mathbf{r} = (r_1, \dots, r_n)$, $r_i \in \mathbb{Z}_{>0}$. Define $P_{\mathbf{r}}(x)$ as

$$P_{\mathbf{r}}(x) = P(x^{r_1}, \dots, x^{r_n}),$$

and let

$$q(\mathbf{r}) = \min \left\{ H(\mathbf{s}) : \mathbf{s} = (s_1, \dots, s_n) \in \mathbb{Z}^n, \mathbf{s} \neq (0, \dots, 0), \sum_{j=1}^n s_j r_j = 0 \right\},$$

Then

$$\lim_{q(\mathbf{r}) \rightarrow \infty} m(P_{1\mathbf{r}}, \dots, P_{l\mathbf{r}}) = m(P_1, \dots, P_l).$$

- $$\lim_{n \rightarrow \infty} m_{2h+1} \left(\frac{x^n - 1}{x - 1} \right) = m_{2h+1} \left(\frac{y - 1}{x - 1} \right) = 0.$$

- $$m_{2h+1} \left(\frac{x^n - 1}{x - 1} \right) = C(h) \frac{\log^{2h-1} n}{n} \left(1 + O \left(\log^{-1} n \right) \right).$$

(suggested by Soundararajan)

Summing up

- Lehmer's question FALSE for m_{2h} , $h \geq 1$.
- Lehmer's question TRUE for m_{2h+1} , $h \geq 1$.
- $m_k(P)$ is interesting for P cyclotomic and that answers Lehmer's question.

Further questions

- $m_3(P)$ for P noncyclotomic?
- Lehmer's question for P noncyclotomic.
- Bounds for $m_{2h}(P)$, $h \geq 2$.
- Explain zeta values!!!

Thank you!
Merci!