

Some aspects of Mahler Measure

Number Theory Seminar – Texas A & M University

March 10th, 2005

Matilde N. Lalin

1. Mahler measure

Definition 1 For $P \in \mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$, the (logarithmic) Mahler measure is defined by

$$m(P) = \frac{1}{(2\pi i)^n} \int_{\mathbb{T}^n} \log |P(x_1, \dots, x_n)| \frac{dx_1}{x_1} \dots \frac{dx_n}{x_n}. \quad (1)$$

This integral is not singular and $m(P)$ always exists.

Because of Jensen's formula:

$$\int_0^1 \log |e^{2\pi i \theta} - \alpha| d\theta = \log^+ |\alpha|, \quad (2)$$

¹we have a simple expression for the Mahler measure of one-variable polynomials:

$$m(P) = \log |a_d| + \sum_{n=1}^d \log^+ |\alpha_n| \quad \text{for} \quad P(x) = a_d \prod_{n=1}^d (x - \alpha_n).$$

2. Examples of Mahler measures in several variables

For two and three variables, several examples are known. The first and simplest examples in two and three variables were given by Smyth [19] and also [1]:

$$m(1 + x + y) = \frac{1}{\pi} D(\zeta_6) = \frac{3\sqrt{3}}{4\pi} L(\chi_{-3}, 2) = L'(\chi_{-3}, -1) \quad (3)$$

$$m(1 + x + y + z) = \frac{7}{2\pi^2} \zeta(3) \quad (4)$$

Other kinds of examples are the families studied by Boyd–Rodriguez-Villegas [18]

$$m\left(x + \frac{1}{x} + y + \frac{1}{y} - k\right) \stackrel{?}{=} \frac{L'(E_k, 0)}{B_k} \quad k \in \mathbb{N}$$

$$m\left(x + \frac{1}{x} + y + \frac{1}{y} - 4\right) = 2L'(\chi_{-4}, -1)$$

$$m\left(x + \frac{1}{x} + y + \frac{1}{y} - 4\sqrt{2}\right) = L'(A, 0)$$

¹ $\log^+ x = \log \max\{1, x\}$ for $x \in \mathbb{R}_{\geq 0}$

Where B_k is a rational number, and E_k is the elliptic curve with corresponds to the zero set of the polynomial. When $k = 4$ the curve has genus zero. When $k = 4\sqrt{2}$ the elliptic curve is

$$A : y^2 = x^3 - 44x + 112,$$

which has complex multiplication.

3. Polylogarithms

Many examples should be understood in the context of polylogarithms.

Definition 2 *The k th polylogarithm is the function defined by the power series*

$$\text{Li}_k(x) := \sum_{n=1}^{\infty} \frac{x^n}{n^k} \quad x \in \mathbb{C}, \quad |x| < 1. \quad (5)$$

This function can be continued analytically to $\mathbb{C} \setminus [1, \infty)$.

In order to avoid discontinuities, and to extend polylogarithms to the whole complex plane, several modifications have been proposed. Zagier [22] considers the following version:

$$P_k(x) := \text{Re}_k \left(\sum_{j=0}^k \frac{2^j B_j}{j!} (\log |x|)^j \text{Li}_{k-j}(x) \right), \quad (6)$$

where B_j is the j th Bernoulli number, $\text{Li}_0(x) \equiv -\frac{1}{2}$ and Re_k denotes Re or Im depending on whether k is odd or even.

This function is one-valued, real analytic in $\mathbb{P}^1(\mathbb{C}) \setminus \{0, 1, \infty\}$ and continuous in $\mathbb{P}^1(\mathbb{C})$. Moreover, P_k satisfy very clean functional equations. The simplest ones are

$$P_k \left(\frac{1}{x} \right) = (-1)^{k-1} P_k(x) \quad P_k(\bar{x}) = (-1)^{k-1} P_k(x).$$

There are also lots of functional equations which depend on the index k . For instance, for $k = 2$, we have the Bloch–Wigner dilogarithm,

$$D(x) := \text{Im}(\text{Li}_2(x)) + \arg(1-x) \log |x|$$

which satisfies the well-known five-term relation

$$D(x) + D(1-xy) + D(y) + D\left(\frac{1-y}{1-xy}\right) + D\left(\frac{1-x}{1-xy}\right) = 0. \quad (7)$$

4. Mahler measure and hyperbolic volumes

A generalization of Smyth's first result was due to Cassaigne and Maillot [17]: for $a, b, c \in \mathbb{C}^*$,

$$\pi m(a + bx + cy) = \begin{cases} D\left(\left|\frac{a}{b}\right| e^{i\gamma}\right) + \alpha \log |a| + \beta \log |b| + \gamma \log |c| & \Delta \\ \pi \log \max\{|a|, |b|, |c|\} & \text{not } \Delta \end{cases} \quad (8)$$

where Δ stands for the statement that $|a|$, $|b|$, and $|c|$ are the lengths of the sides of a triangle, and α , β , and γ are the angles opposite to the sides of lengths $|a|$, $|b|$, and $|c|$




Figure 1: The main term in Cassaigne – Maillot formula is the volume of the ideal hyperbolic tetrahedron over the triangle.

respectively. The term with the dilogarithm can be interpreted as the volume of the ideal hyperbolic tetrahedron which has the triangle as basis and the fourth vertex is infinity. See figure 1.

Another example was due to Vandervelde [20]. He studied the polynomials whose equation can be expressed as

$$y = \frac{bx + d}{ax + c}.$$

When $a, b, c, d \in \mathbb{R}^*$, the Mahler measure of this polynomial is the sum of some logarithms and two dilogarithm terms, which can be interpreted as the volume of the ideal polyhedra built over a cyclic quadrilateral of sides $|a|$, $|b|$, $|c|$ and $|d|$.

We have studied the case of

$$y = \frac{x^n - 1}{t(x^m - 1)} = \frac{x^{n-1} + \dots + 1}{t(x^{m-1} + \dots + 1)}$$

and obtained a similar result, the Mahler measure is given by a formula whose dilogarithm terms are the volumes of ideal polyhedra that are constructed over all the possible polygons with m sides of length $|t|$ and n sides of length 1.

Moreover, this phenomenon is similar to the A -polynomial phenomenon described by Boyd [3] and Boyd and Rodriguez Villegas [5] as we showed that this polynomial can be thought as an analogous for an A -polynomial. More specifically, we showed that it may be obtained a factor of the resultant of certain gluing and completeness equations (conveniently modified by the deformation parameters) in the similar way as A -polynomials are obtained.

5. More examples of Mahler measures in several variables

For more than three variables, very little is known.

Theorem 3 For $n \geq 1$ we have:

$$\begin{aligned} & \pi^{2n} m \left(1 + \left(\frac{1-x_1}{1+x_1} \right) \dots \left(\frac{1-x_{2n}}{1+x_{2n}} \right) z \right) \\ &= \sum_{h=1}^n \frac{s_{n-h}(2^2, \dots, (2n-2)^2)}{(2n-1)!} \pi^{2n-2h} (2h)! \frac{2^{2h+1} - 1}{2} \zeta(2h+1) \end{aligned} \tag{9}$$

For $n \geq 0$:

$$\begin{aligned} & \pi^{2n+1} m \left(1 + \left(\frac{1-x_1}{1+x_1} \right) \cdots \left(\frac{1-x_{2n+1}}{1+x_{2n+1}} \right) z \right) \\ &= \sum_{h=0}^n \frac{s_{n-h}(1^2, \dots, (2n-1)^2)}{(2n)!} 2^{2h+1} \pi^{2n-2h} (2h+1)! \mathbf{L}(\chi_{-4}, 2h+2) \end{aligned} \quad (10)$$

There are analogous (but more complicated) formulas for

$$\begin{aligned} & m \left(1 + x + \left(\frac{1-x_1}{1+x_1} \right) \cdots \left(\frac{1-x_n}{1+x_n} \right) (1+y)z \right) \\ & m \left(1 + \left(\frac{1-x_1}{1+x_1} \right) \cdots \left(\frac{1-x_n}{1+x_n} \right) x + \left(1 - \left(\frac{1-x_1}{1+x_1} \right) \cdots \left(\frac{1-x_n}{1+x_n} \right) \right) y \right) \end{aligned}$$

Where

$$s_l(a_1, \dots, a_k) = \begin{cases} 1 & \text{if } l = 0 \\ \sum_{i_1 < \dots < i_l} a_{i_1} \cdots a_{i_l} & \text{if } 0 < l \leq k \\ 0 & \text{if } k < l \end{cases} \quad (11)$$

are the elementary symmetric polynomials, i. e.,

$$\prod_{i=1}^k (x + a_i) = \sum_{l=0}^k s_l(a_1, \dots, a_k) x^{k-l} \quad (12)$$

For example,

$$\pi^3 m \left(1 + \left(\frac{1-x_1}{1+x_1} \right) \left(\frac{1-x_2}{1+x_2} \right) \left(\frac{1-x_3}{1+x_3} \right) z \right) = 24\mathbf{L}(\chi_{-4}, 4) + \pi^2 \mathbf{L}(\chi_{-4}, 2) \quad (13)$$

$$\pi^4 m \left(1 + \left(\frac{1-x_1}{1+x_1} \right) \cdots \left(\frac{1-x_4}{1+x_4} \right) z \right) = 62\zeta(5) + \frac{14\pi^2}{3}\zeta(3) \quad (14)$$

$$\pi^4 m \left(1 + x + \left(\frac{1-x_1}{1+x_1} \right) \left(\frac{1-x_2}{1+x_2} \right) (1+y)z \right) = 93\zeta(5) \quad (15)$$

$$(16)$$

6. Examples coming from the world of resultants

Let us mention some examples of Mahler measure of resultants (this will be part of a joint work with D'Andrea [7]).

Theorem 4

$$\begin{aligned} m(\text{Res}_{\{0,m,n\}}) &= m(\text{Res}_t(x + yt^m + t^n, z + wt^m + t^n)) = m \left(z - \frac{(1-x)^m(1-y)^{n-m}}{(1-xy)^n} \right) \\ &= \frac{2}{\pi^2} (-mP_3(\varphi^n) - nP_3(-\varphi^m) + mP_3(\phi^n) + nP_3(\phi^m)) \end{aligned}$$

where φ is the real root of $x^n + x^{n-m} - 1 = 0$ such that $0 \leq \varphi \leq 1$, and ϕ is the real root of $x^n - x^{n-m} - 1 = 0$ such that $1 \leq \phi$. In particular, for $m = 1, n = 2$,

$$m(P) = \frac{4}{\pi^2} (P_3(\phi) - P_3(-\phi)) \quad (17)$$

where $\phi^2 + \phi - 1 = 0$ and $0 \leq \phi \leq 1$ (in other words, $\phi = \frac{-1+\sqrt{5}}{2}$). Moreover, using the numerical identity

$$\frac{\zeta_{\mathbb{Q}(\sqrt{5})}(3)}{\zeta(3)} \stackrel{?}{=} \frac{1}{\sqrt{5}}(P_3(\phi) - P_3(-\phi))$$

(see Zagier [21]), then

$$m(\text{Res}_{\{0,1,2\}}) \stackrel{?}{=} \frac{4\sqrt{5}\zeta_{\mathbb{Q}(\sqrt{5})}(3)}{\pi^2\zeta(3)}$$

Theorem 5

$$\begin{aligned} m(\text{Res}_{\{(0,0),(1,0),(0,1)\}}) &= m\left(\begin{array}{ccc|c} x & y & z & \\ u & v & w & \\ r & s & t & \end{array}\right) \\ &= m((1-x)(1-y) - (1-z)(1-w)) = \frac{9\zeta(3)}{2\pi^2} \end{aligned}$$

7. Beilinson's conjectures

One of the main problems in Number Theory is finding rational (or integral) solutions of polynomial equations with rational coefficients (global solutions). In spite of the failure of the local-global principle in general, there are several theorems and conjectures which predict that one may obtain global information from local information and that that relation is made through values of L-functions. These statements include the Dirichlet class number formula, the Birch–Swinnerton-Dyer conjecture, and more generally, Bloch's and Beilinson's conjectures.

Typically, there are four elements involved in this setting: an arithmetic-geometric object X (typically, an algebraic variety), its L-function (which codify local information), a finitely generated abelian group K , and a regulator map $K \rightarrow \mathbb{R}$. When K has rank 1, Beilinson's conjectures predict that the $L'_X(0)$ is, up to a rational number, equal to a value of the regulator.

For instance, for a number field F , Dirichlet class number formula states that

$$\lim_{s \rightarrow 1} (s-1)\zeta_F(s) = \frac{2^{r_1}(2\pi)^{r_2} h_F \text{reg}_F}{\omega_F \sqrt{|D_F|}}.$$

Here, $X = \mathcal{O}_F$ (the ring of integers), $L_X = \zeta_F$, and the group is \mathcal{O}_F^* . Hence, when F is a real quadratic field, Dirichlet class number formula may be written as $\zeta'_F(0)$ is equal to, up to a rational number, $\log |\epsilon|$, for some $\epsilon \in \mathcal{O}_F^*$.

8. An algebraic integration for Mahler measure

The appearance of L-functions in Mahler measures formulas is a common phenomenon. Deninger [8] interpreted the Mahler measure as a Deligne period of a mixed motive. More specifically, in two variables, and under certain conditions, he proved that

$$m(P) = \text{reg}(\xi_i),$$

where reg is the determinant of the regulator matrix, which we are evaluating in some class in an appropriate group in K -theory.

Rodriguez-Villegas [18] has worked out the details for two variables. This was further developed by Boyd and Rodriguez-Villegas [4], [5].

More specifically one has

$$m(P) = m(P^*) - \frac{1}{2\pi} \int_{\gamma} \eta(x, y), \quad (18)$$

where

$$\eta(x, y) = \log |x| d \arg y - \log |y| d \arg x \quad (19)$$

is a differential form that is "essentially" defined in the curve C determined by the zeros of P . This form is essentially the regulator.

One has a crucial property:

Theorem 6

$$\eta(x, 1-x) = dD(x). \quad (20)$$

Because of the above property, there is a condition that tells us when $\eta(x, y)$ is exact, namely:

$$x \wedge y = \sum_j r_j z_j \wedge (1 - z_j)$$

in $\wedge^2(\mathbb{C}(C)^*) \otimes \mathbb{Q}$, in other words, $\{x, y\} = 0$ in $K_2(\mathbb{C}(C)) \otimes \mathbb{Q}$.

Under those circumstances,

$$\eta(x, y) = d \left(\sum_j r_j D(z_j) \right) = dD \left(\sum_j r_j [z_j] \right).$$

We have $\gamma \subset C$ such that

$$\partial\gamma = \sum_k \epsilon_k [w_k] \quad \epsilon_k = \pm 1$$

where $w_k \in C(\mathbb{C})$, $|x(w_k)| = |y(w_k)| = 1$. Then

$$2\pi m(P) = D(\xi) \quad \text{for } \xi = \sum_k \sum_j r_j [z_j(w_k)].$$

We could summarize the whole picture as follows:

$$\dots \rightarrow (K_3(\bar{\mathbb{Q}}) \supset) K_3(\partial\gamma) \rightarrow K_2(C, \partial\gamma) \rightarrow K_2(C) \rightarrow \dots$$

$$\partial\gamma = C \cap \mathbb{T}^2$$

There are two "nice" situations:

- $\eta(x, y)$ is exact, then $\{x, y\} \in K_3(\partial\gamma)$. In this case we have $\partial\gamma \neq \emptyset$, we use Stokes' Theorem and we finish with an element $K_3(\partial\gamma) \subset K_3(\bar{\mathbb{Q}})$, leading to dilogarithms and zeta functions (of number fields), due to theorems by Borel, Bloch, Suslin and others.
- $\partial\gamma = \emptyset$, then $\{x, y\} \in K_2(C)$. In this case, we have $\eta(x, y)$ is not exact and we get essentially the L -series of a curve, leading to examples of Beilinson's conjectures.

In general, we may get combinations of both situations.

9. The three-variable case

We are going to extend this situation to three variables. We will take

$$\begin{aligned} \eta(x, y, z) &= \log |x| \left(\frac{1}{3} d \log |y| d \log |z| - d \arg y d \arg z \right) \\ &+ \log |y| \left(\frac{1}{3} d \log |z| d \log |x| - d \arg z d \arg x \right) + \log |z| \left(\frac{1}{3} d \log |x| d \log |y| - d \arg x d \arg y \right) \end{aligned}$$

Then η verifies

$$d\eta(x, y, z) = \operatorname{Re} \left(\frac{dx}{x} \wedge \frac{dy}{y} \wedge \frac{dz}{z} \right),$$

so it is closed.

We can express the Mahler measure of P

$$m(P) = m(P^*) - \frac{1}{(2\pi)^2} \int_{\Gamma} \eta(x, y, z).$$

Where

$$\Gamma = \{P(x, y, z) = 0\} \cap \{|x| = |y| = 1, |z| \geq 1\}.$$

We are integrating on a subset of $S = \{P(x, y, z) = 0\}$. The differential form is defined in this surface minus the set of zeros and poles of x , y and z , but that will not interfere our purposes, since we will be dealing with the cases when $\eta(x, y, z)$ is exact and that implies trivial tame symbols thus the element in the cohomology can be extended to S .

As in the two-variable case, we would like to apply Stokes' Theorem.

Let us take a look at Smyth's case, we can express the polynomial as $P(x, y, z) = (1 - x) + (1 - y)z$. We get:

$$m(P) = m(1 - y) + \frac{1}{(2\pi i)^2} \int_{\mathbb{T}^2} \log^+ \left| \frac{1 - x}{1 - y} \right| \frac{dx}{x} \frac{dy}{y} = -\frac{1}{(2\pi)^2} \int_{\Gamma} \eta(x, y, z).$$

In general, we have

$$\eta(x, 1 - x, y) = d\omega(x, y),$$

where

$$\omega(x, y) = -D(x) d \arg y + \frac{1}{3} \log |y| (\log |1 - x| d \log |x| - \log |x| d \log |1 - x|).$$

Suppose we have

$$x \wedge y \wedge z = \sum r_i x_i \wedge (1 - x_i) \wedge y_i$$

in $\wedge^3(\mathbb{C}(S)^*) \otimes \mathbb{Q}$.

Then

$$\int_{\Gamma} \eta(x, y, z) = \sum r_i \int_{\Gamma} \eta(x_i, 1 - x_i, y_i) = \sum r_i \int_{\partial\Gamma} \omega(x_i, y_i).$$

In Smyth's case, this corresponds to

$$x \wedge y \wedge z = -x \wedge (1 - x) \wedge y - y \wedge (1 - y) \wedge x,$$

in other words,

$$\eta(x, y, z) = -\eta(x, 1-x, y) - \eta(y, 1-y, x).$$

Back to the general picture, $\partial\Gamma = \{P(x, y, z) = 0\} \cap \{|x| = |y| = |z| = 1\}$. When $P \in \mathbb{Q}[x, y, z]$, Γ can be thought as

$$\gamma = \{P(x, y, z) = P(x^{-1}, y^{-1}, z^{-1}) = 0\} \cap \{|x| = |y| = 1\}.$$

Note that we are integrating now on a path inside the curve $C = \{P(x, y, z) = P(x^{-1}, y^{-1}, z^{-1}) = 0\}$. The differential form ω is defined in this new curve (this way of thinking the integral over a new curve has been proposed by Maillot). Now it makes sense to try to apply Stokes' Theorem again. We have

$$\omega(x, x) = dP_3(x).$$

Suppose we have

$$[x]_2 \otimes y = \sum r_i [x_i]_2 \otimes x_i$$

in $(B_2(\mathbb{C}(C)) \otimes \mathbb{C}(C)^*)_{\mathbb{Q}}$.


Then, as before:

$$\int_{\gamma} \omega(x, y) = \sum r_i P_3(x_i)|_{\partial\gamma}.$$

Back to Smyth's case, in order to compute C we set $\frac{(1-x)(1-x^{-1})}{(1-y)(1-y^{-1})} = 1$ and we get $C = \{x = y\} \cup \{xy = 1\}$ in this example, and

$$-[x]_2 \otimes y - [y]_2 \otimes x = \pm 2[x]_2 \otimes x.$$

We integrate in the set described by the following picture



Then

$$m((1-x) + (1-y)z) = \frac{1}{4\pi^2} \int_{\gamma} \omega(x, y) + \omega(y, x) = \frac{1}{4\pi^2} 8(P_3(1) - P_3(-1)) = \frac{7}{2\pi^2} \zeta(3).$$

10. The K -theory conditions

We follow Goncharov, [9], [10]. Given a field F , we define subgroups $R_i(F) \subset \mathbb{Z}[\mathbb{P}_F^1]$ as

$$R_1(F) := [x] + [y] - [xy]$$

$$R_2(F) := [x] + [y] + [1-xy] + \left[\frac{1-x}{1-xy} \right] + \left[\frac{1-y}{1-xy} \right]$$

$$R_3(F) := \text{certain functional equation of the trilogarithm}$$

Define

$$B_i(F) := \mathbb{Z}[\mathbb{P}_F^1]/R_i(F) \quad (21)$$

The idea is that $B_i(F)$ is the place where P_i naturally acts. We have the following complexes:

$$\begin{aligned} B_F(3) &: B_3(F) \xrightarrow{\delta_1^3} B_2(F) \otimes F^* \xrightarrow{\delta_2^3} \wedge^3 F^* \\ B_F(2) &: B_2(F) \xrightarrow{\delta_1^2} \wedge^2 F^* \\ B_F(1) &: F^* \end{aligned}$$

($B_i(F)$ is placed in degree 1).

$$\delta_1^3([x]_3) = [x]_2 \otimes x \quad \delta_2^3([x]_2 \otimes y) = x \wedge (1-x) \wedge y \quad \delta_1^2([x]_2) = x \wedge (1-x)$$

Proposition 7

$$H^1(B_F(1)) \cong K_1(F) \quad (22)$$

$$H^1(B_F(2))_{\mathbb{Q}} \cong K_3^{\text{ind}}(F)_{\mathbb{Q}} \quad (23)$$

$$H^2(B_F(2)) \cong K_2(F) \quad (24)$$

$$H^3(B_F(3)) \cong K_3^M(F) \quad (25)$$

Goncharov [9] conjectures:

$$H^i(B_F(3) \otimes \mathbb{Q}) \cong K_{6-i}^{[3-i]}(F)_{\mathbb{Q}}$$

Where $K_n^{[i]}(F)_{\mathbb{Q}}$ is a certain quotient in a filtration of $K_n(F)_{\mathbb{Q}}$.

Note that our first condition is that

$$x \wedge y \wedge z = 0 \quad \text{in} \quad H^3(B_{\mathbb{Q}(S)}(3) \otimes \mathbb{Q}) \cong K_3^{[0]}(\mathbb{Q}(S))_{\mathbb{Q}} \cong K_3^M(\mathbb{Q}(S)) \otimes \mathbb{Q}$$

and the second condition is

$$[x_i]_2 \otimes y_i = 0 \quad \text{in} \quad H^2(B_{\mathbb{Q}(C)}(3) \otimes \mathbb{Q}) \stackrel{?}{\cong} K_4^{[1]}(\mathbb{Q}(C))_{\mathbb{Q}}$$

Hence, the conditions can be translated as certain elements in different K -theories must be zero, which is analogous to the two-variable case.

We could summarize this picture as follows. We first integrate in this picture

$$\dots \rightarrow K_4(\partial\Gamma) \rightarrow K_3(S, \partial\Gamma) \rightarrow K_3(S) \rightarrow \dots$$

$$\partial\Gamma = S \cap \mathbb{T}^3$$

As before, we have two situations. All the examples we have talked about fit into the situation when $\eta(x, y, z)$ is exact and $\partial\Gamma \neq \emptyset$. Then we finish with an element in $K_4(\partial\Gamma)$.

Then we go to


$$\dots \rightarrow (K_5(\bar{\mathbb{Q}}) \supset) K_5(\partial\gamma) \rightarrow K_4(C, \partial\gamma) \rightarrow K_4(C) \rightarrow \dots$$

$$\partial\gamma = C \cap \mathbb{T}^2$$

Again we have two possibilities, but in our context, $\omega(x, y)$ is exact and we finish with an element in $K_5(\partial\gamma) \subset K_5(\overline{\mathbb{Q}})$ leading to trilogarithms and zeta functions, due to Zagier's conjecture and Borel's theorem.

11. Conclusion

The next picture shows how Mahler measure interacts with several elements (some have been discussed here and some have not). We can see the key role of Mahler measure in the relation among special values of L-functions and regulators (which are related via Beilinson's conjectures), heights, and hyperbolic manifolds (that are related by Beilinson's conjectures as well). It is our general goal to bring more light to the nature of these relationships.



References

- [1] D. W. Boyd, Speculations concerning the range of Mahler's measure, *Canad. Math. Bull.* **24** (1981), 453 - 469.
- [2] D. W. Boyd, Mahler's measure and special values of L-functions, *Experiment. Math.* **7** (1998), 37 - 82.
- [3] D. W. Boyd, Mahler's measure and invariants of hyperbolic manifolds, *Number Theory for the Millennium* (M.A. Bennett et al., eds.), A K Peters, Boston (2002) pp. 127 -143.
- [4] D. W. Boyd, F. Rodriguez Villegas, Mahler's measure and the dilogarithm (I), *Canad. J. Math.* **54** (2002), 468 - 492.
- [5] D. W. Boyd, F. Rodriguez Villegas, with an appendix by N. M. Dunfield, Mahler's measure and the dilogarithm (II), (preprint, July 2003)
- [6] J. Condon, Calculation of the Mahler measure of a three variable polynomial, (preprint, October 2003)
- [7] C. D'Andrea, M. N. Lalín, On The Mahler measure of resultants in small dimensions. (in preparation).
- [8] C. Deninger, Deligne periods of mixed motives, *K-theory and the entropy of certain Z^n -actions*, *J. Amer. Math. Soc.* **10** (1997), no. 2, 259-281.

- [9] A. B. Goncharov, Geometry of Configurations, Polylogarithms, and Motivic Cohomology, *Adv. Math.* **114** (1995), no. 2, 197–318.
- [10] A. B. Goncharov, The Classical Polylogarithms, Algebraic K-theory and $\zeta_F(n)$, *The Gelfand Mathematical Seminars 1990-1992*, Birkhauser, Boston (1993), 113 - 135.
- [11] R. Kellerhals, On the volumes of hyperbolic 5-orthoschemes and the trilogarithm, *Comment. Math. Helv.* **67** (1992), no. 4, 648–663.
- [12] R. Kellerhals, Volumes in hyperbolic 5-space. *Geom. Funct. Anal.* **5** (1995), no. 4, 640–667.
- [13] M. N. Lalín, Some examples of Mahler measures as multiple polylogarithms, *J. Number Theory* **103** (2003), no. 1, 85–108.
- [14] M. N. Lalín, Mahler Measure and Volumes in Hyperbolic Space, (September 2003), to appear in *Geom. Dedicata*
- [15] M. N. Lalín, Mahler measure of some n-variable polynomial families, (September 2004), submitted to *J. Number Theory*
- [16] M. N. Lalín, An algebraic integration for Mahler measure (in preparation).
- [17] V. Maillot, Géométrie d’Arakelov des variétés toriques et fibrés en droites intégrables. *Mém. Soc. Math. Fr. (N.S.)* **80** (2000), 129pp.
- [18] F. Rodriguez Villegas, Modular Mahler measures I, *Topics in number theory* (University Park, PA 1997), 17-48, Math. Appl., 467, Kluwer Acad. Publ. Dordrecht, 1999.
- [19] C. J. Smyth, On measures of polynomials in several variables, *Bull. Austral. Math. Soc. Ser. A* **23** (1981), 49 - 63. Corrigendum (with G. Myerson): *Bull. Austral. Math. Soc.* **26** (1982), 317 - 319.
- [20] S. Vandervelde, A formula for the Mahler measure of $axy + bx + cy + d$, *J. Number Theory* **100** (2003) 184–202.
- [21] D. Zagier, The Bloch-Wigner-Ramakrishnan polylogarithm function. *Math. Ann.* **286** (1990), no. 1-3, 613–624.
- [22] D. Zagier, Polylogarithms, Dedekind Zeta functions, and the Algebraic K-theory of Fields, *Arithmetic algebraic geometry* (Texel, 1989), 391–430, Progr. Math., 89, Birkhuser Boston, Boston, MA, 1991.