FAILURE OF THE WELL-ROUNDED RETRACT FOR OUTER SPACE AND TEICHMÜLLER SPACE

MAXIME FORTIER BOURQUE

ABSTRACT. The well-rounded retract for $SL_n(\mathbb{Z})$ is defined as the set of flat tori of unit volume and dimension n whose systoles generate a finite-index subgroup in homology. This set forms an equivariant spine of minimal dimension for the space of flat tori.

For both the Outer space X_n of metric graphs of rank n and the Teichmüller space \mathcal{T}_g of closed hyperbolic surfaces of genus g, we show that the literal analogue of the well-rounded retract does not contain an equivariant spine. We also prove that the sets of graphs whose systoles fill either topologically or geometrically (two analogues of a set proposed as a spine for \mathcal{T}_g by Thurston) are spines for X_n but that their dimension is larger than the virtual cohomological dimension of $Out(F_n)$ in general.

1. INTRODUCTION

A systole in a compact metric space is a non-contractible closed curve of minimal length among such curves. Ash [Ash77] defined the well-rounded retract \mathfrak{W}_n of the space \mathfrak{T}_n of marked flat tori of unit volume and dimension n as the set of tori T whose systoles generate a finite-index subgroup in $H_1(T,\mathbb{Z})$, following work of Soulé [Sou75] in the case n = 3. He proved that there is an $\mathrm{SL}_n(\mathbb{Z})$ -equivariant deformation retraction of \mathfrak{T}_n onto \mathfrak{W}_n , i.e., that \mathfrak{W}_n is an equivariant spine for \mathfrak{T}_n . Furthermore, the quotient $\mathfrak{W}_n/\mathrm{SL}_n(\mathbb{Z})$ is compact and the dimension of \mathfrak{W}_n is equal to the virtual cohomological dimension (vcd) of $\mathrm{SL}_n(\mathbb{Z})$, the smallest possible for a spine.

Motivated by this, Thurston [Thu85] considered the set \mathcal{V}_g of marked closed hyperbolic surfaces of genus g whose systoles *fill*, meaning that each component of the complement of their union is contractible (hence the interior of a polygon). Equivalently, a set C of curves on a surface S fills if every non-contractible closed curve in S intersects some element of C. Thurston sketched a proof that there is a mapping class group equivariant deformation retract of the Teichmüller space \mathcal{T}_g onto \mathcal{V}_g , but his argument had gaps [Ji14]. Furthermore, the dimension of \mathcal{V}_g is larger than the vcd of the mapping class group Mod_g in general [FB23].

The third character in this story is the Culler–Vogtmann Outer space X_n of marked metric graphs of unit volume and rank equal to n, upon which the group $Out(F_n)$ of outer automorphisms of the free group of rank n acts. Culler and Vogtmann [CV86] found a cocompact equivariant spine K_n for

 X_n of dimension 2n-3, equal to the vcd of $Out(F_n)$ (which was determined using K_n). This spine K_n is not defined in terms of systeles.

If (E, G) is equal to either $(\mathfrak{T}_n, \mathrm{SL}_n(\mathbb{Z})), (\mathcal{T}_g, \mathrm{Mod}_g)$, or $(X_n, \mathrm{Out}(F_n))$, and $x \in E$, then

$$\dim(E) = \operatorname{vcd}(G) + \operatorname{rank}(H_1(x,\mathbb{Z})) - 1,$$

which suggests that one should use homology to define spines of minimal dimension. A naive approach is to simply transpose the definition of the well-rounded retract \mathfrak{W}_n in the other two settings. That is, we can define the set $W_n \subset X_n$ of graphs whose systoles generate a finite-index subgroup in integral homology and the set $\mathcal{W}_g \subset \mathcal{T}_g$ of hyperbolic surfaces whose systoles generate a finite-index subgroup in integral homology. In his PhD thesis [Bak11], Baker proved that W_3 is a spine of minimal dimension for X_3 different from K_3 . However, these analogues W_n and \mathcal{W}_g of the well-rounded retract fail to achieve their goal in general.

Theorem 1.1. There exist infinitely many $n \ge 2$ such that W_n does not contain any $Out(F_n)$ -equivariant spine for X_n .

Theorem 1.2. There exist infinitely many $g \ge 2$ such that \mathcal{W}_g does not contain any Mod_q -equivariant spine for \mathcal{T}_q .

Note that the dimension of W_n (resp. W_g) is equal to the vcd of $Out(F_n)$ (resp. Mod_g). The obstruction comes instead from the fact that these sets miss certain loci of fixed points of finite subgroups that have to intersect any spine.

These theorems go in the same direction as results of Pettet and Souto showing that \mathfrak{W}_n is a minimal spine [PS08a] and slightly modifying its definition can yield sets of the same dimension that are not spines anymore [PS08b]. In other words, spines are sensitive and thus tricky to find.

There is also an analogue of the Thurston set \mathcal{V}_g in X_n . Indeed, consider the set $V_n \subset X_n$ of graphs whose systoles *topologically fill*, meaning that each component of the complement of their union is contractible. Equivalently, a set C of closed geodesics in a metric graph Γ topologically fills if every non-contractible curve in Γ intersects some element of C. One could also consider the set V'_n of graphs whose systoles geometrically fill in the sense that their union is equal to the whole graph. It is easy to see that

$$W_n \subseteq V_n$$
 and $V'_n \subseteq V_n$

Furthermore, V'_2 coincides with K_2 , the dual to the Farey triangulation, but $W_2 = V_2$ is strictly larger (it contains the dumbbells with two loops of equal length, which form spikes emanating from the midpoints of the edges in K_2).

In contrast with W_n , the sets V_n and V'_n are always spines.

Theorem 1.3. For every $n \ge 2$, the set V_n is an $Out(F_n)$ -equivariant spine for X_n and V'_n is an equivariant spine for V_n .

However, their dimension is too large in general.

Theorem 1.4. For every $\varepsilon \in (0,1)$, there exists an n such that the dimension of V'_n is at least $(3 - \varepsilon)n$, hence larger than the vcd of $\operatorname{Out}(F_n)$.

Note that there is also a dynamically-defined notion of filling currents for free groups due to Kapovich and Lustig [KL10]. We do not know whether the set of graphs whose systoles fill in that sense forms a spine and if so, what its dimension is.

One may wonder if there is a spine of the minimal dimension 2n-3 contained in V'_n . However, it seems difficult to push the deformation retraction defined in the proof of Theorem 1.3 much further. One can continue until there is a systole passing through any pair of edges that are adjacent at a vertex of degree 3 by folding these edges gradually otherwise, but the proof of Theorem 1.4 implies that the dimension of the resulting set is still too large in general.

Acknowledgements. I thank Thomas Haettel for making a remark which sparked this project after a talk I gave at UQAM, Karen Vogtmann for pointing out the reference [Bak11], and the referee for useful comments and corrections.

2. Outer space

We start by proving the negative results regarding Outer space. The proof of both Theorem 1.1 and Theorem 1.4 is based on the same family of graphs that have a large automorphism group and few systoles that cover the whole graph. These graphs were used in [FB20] to construct hyperbolic surfaces with similar properties.

Given integers $p, q \geq 2$, a map of type $\{p, q\}$ is a connected graph of constant valence (degree) q embedded in an oriented surface such that each complementary region (whose closure is called a *face*) is a topological disk whose boundary consists of p edges. This can also be phrased in terms of a ribbon structure on the graph. A *flag* is a triple (v, e, f) where v is a vertex, e is an edge, f is a face, and $v \subset e \subset f$. A map is *flag-transitive* if for any two flags there is a homeomorphism of the underlying surface which sends the map to itself and the first flag to the second. For now we consider our maps as combinatorial graphs where each edge has length 1. The *girth* of a combinatorial graph is the same as its systole, namely, the minimal length of a cycle that is not contractible.

We will require a small variation of a result of Evans [Eva79, Theorem 11] about the existence of flag-transitive maps of large girth. The difference here is that we want to make sure that only the obvious cycles have length equal to the girth.

Lemma 2.1. For any $q \ge 3$ and $p \ge 7$, there exists a finite flag-transitive map M of type $\{p,q\}$ and girth p such that the only non-trivial cycles of length p in M are the face boundaries.

Proof. There is an infinite flag-transitive map $M_{p,q}$ of type $\{p,q\}$ embedded in the hyperbolic plane \mathbb{H}^2 coming from the tiling by regular *p*-gons with interior angles $2\pi/q$. The automorphisms of $M_{p,q}$ are realized by a finitely-generated discrete group *G* of isometries of the hyperbolic plane. By Mal'cev's theorem [Mal65], *G* is residually finite, so there is a sequence of normal subgroups $G_k \triangleleft G$ of finite index such that $\bigcap G_k = \{\text{id}\}$. This implies that G_k is eventually torsion-free and the closed hyperbolic surfaces $S_k = \mathbb{H}^2/G_k$ have injectivity radius going to infinity as $k \to \infty$. If *k* is large enough, then the projection M_k of $M_{p,q}$ to S_k has type $\{p,q\}$ because the map $\mathbb{H}^2 \to S_k$ is a covering map. Furthermore, M_k is finite since $M_{p,q}/G$ is a half-edge and G_k has finite index in *G*. Lastly, M_k is flag-transitive via the quotient group G/G_k acting on S_k .

Since the face boundaries in M_k have combinatorial length p, the girth of M_k is at most p. Since the injectivity radius of S_k tends to infinity, any cycle in M_k which is not contractible in S_k becomes arbitrarily long (with respect to the hyperbolic metric and therefore also in terms of its number of edges) as k tends to infinity. In particular, a cycle in M_k that is not contractible in S_k has combinatorial length strictly larger than p if k is large enough. It is also true that any cycle in M_k which is contractible in S_k (and hence lifts to the universal cover) has combinatorial length at least p with equality only if it is the boundary of a face. We can prove this as follows. Suppose that γ is an embedded cycle of combinatorial length at most p in $M_{p,q}$. Let A be the hyperbolic area of any face in the tiling, let N be the number of faces enclosed by γ , and for a vertex $v \in \gamma$ let k_v be the geodesic curvature of γ at v, that is, π minus the interior angle. Then the Gauss-Bonnet formula yields

$$2\pi = \sum_{v \in \gamma} k_v - N \cdot A \le p\left(1 - \frac{2}{q}\right)\pi - A = 2\pi$$

so that in fact N = 1 and γ is the boundary of a face.

To prove our results, we use this construction with q = 3 and $p \ge 7$ arbitrarily large. Let M be a map satisfying the conclusions of Lemma 2.1 with these parameters and let V, E, and F be its number of vertices, edges, and faces respectively. Then

$$3V = 2E = pF.$$

The rank n of M is such that its Euler characteristic is

$$1 - n = V - E = -V/2$$

so that n = 1 + V/2. By the lemma, the systoles in M are the faces boundaries, so there are $F = 3V/p = \frac{6}{p}(n-1)$ of them. In particular, the number of systoles divided by the rank n is arbitrarily small if p is large enough.

Proof of Theorem 1.4. Given $\varepsilon \in (0, 1)$, choose $p \geq 7$ such that $6/p < \varepsilon$, then let M be a finite map of type $\{p, 3\}$ as above whose combinatorial systoles are the face boundaries.

Let n be the rank of M, pick an arbitrary homotopy equivalence from the bouquet on n circles to M to get a marking, and make all edges of Mof equal length 1/E so that its volume is 1. We can now consider M as an element in the Outer space X_n . Since the systoles in M are the face boundaries, they cover the whole graph so that $M \in V'_n$.

We now want to deform M (i.e., vary the lengths on its edges) in such a way that the systoles stay the same curves and thus still cover the whole graph. Since competing curves are longer by a definite amount, near Mthese curves will remain systoles as long as they stay of equal length.

If $\gamma_1, \ldots, \gamma_F$ are the systoles, then this requires F - 1 equations, namely,

$$\ell(\gamma_1) = \ell(\gamma_2), \quad \ell(\gamma_2) = \ell(\gamma_3), \quad \dots, \quad \ell(\gamma_{F-1}) = \ell(\gamma_F).$$

In turn, each $\ell(\gamma_j)$ is equal to the sum of the lengths of the edges traversed by γ_j , so this gives us F - 1 linear equations for the edge lengths. The subspace of \mathbb{R}^E cut out by these equations has codimension at most F - 1and then we intersect this with the hyperplane where the sum of the lengths is equal to 1. The dimension of the intersection I is at least

$$E - F = \left(3 - \frac{6}{p}\right)(n-1)$$

and this is larger than $(3-\varepsilon)n$ provided that p (and hence n) is large enough.

As explained above, there is a neighborhood U of M in I where the face boundaries will remain systoles and hence $U \subset V'_n$. This shows that the dimension of V'_n is at least $(3 - \varepsilon)n$. Since $\varepsilon < 1$, this is strictly larger than 2n - 3, the vcd of $Out(F_n)$.

To prove that the well-rounded set W_n does not contain an equivariant spine, we will use the above construction together with the following elementary observation, in which Fix(H) denotes the set of all points fixed by all the elements in H.

Lemma 2.2. Let G be a group acting on a topological space E and let $S \subseteq E$ be a G-equivariant spine for E. If H is a subgroup of G, then $S \cap \text{Fix}(H)$ is a G-equivariant spine for Fix(H). In particular, if $\text{Fix}(H) \neq \emptyset$ then $S \cap \text{Fix}(H) \neq \emptyset$.

Proof. Let $(x, t) \mapsto f_t(x)$ be a continuous map from $E \times [0, 1]$ to E such that f_0 is the identity on E, $f_1(E) = S$, $f_t(x) = x$ for every $x \in S$ and every $t \in [0, 1]$, and $f_t(g(x)) = g(f_t(x))$ for every $g \in G$, every $x \in X$, and every $t \in [0, 1]$. Then for every $t \in [0, 1]$, every $h \in H$, and every $x \in \text{Fix}(H)$, we have

$$h(f_t(x)) = f_t(h(x)) = f_t(x)$$

so that $f_t(x) \in Fix(H)$. This shows that f_t restricts to a map from Fix(H) to Fix(H) for all $t \in [0, 1]$. This restriction is still *G*-equivariant and equal to the identity on $S \cap Fix(H)$. Finally, we have

$$f_1(\operatorname{Fix}(H)) \subseteq f_1(E) \cap \operatorname{Fix}(H) = S \cap \operatorname{Fix}(H) = f_1(S \cap \operatorname{Fix}(H)) \subseteq f_1(\operatorname{Fix}(H))$$

so that $f_1(\operatorname{Fix}(H)) = S \cap \operatorname{Fix}(H)$. In particular, if $\operatorname{Fix}(H) \neq \emptyset$ then

$$S \cap \operatorname{Fix}(H) = f_1(\operatorname{Fix}(H)) \neq \emptyset.$$

We can now prove that W_n does not contain any spine.

Proof of Theorem 1.1. Take any $p \geq 7$ and let M be a finite flag-transitive map of type $\{3, p\}$ such that its systoles are the face boundaries. Recall that there are $\frac{6}{p}(n-1) < (n-1)$ systoles in M where n is the rank. In particular, the systoles in M do not generate a finite-index subgroup in $H_1(M, \mathbb{Z}) \cong \mathbb{Z}^n$. Considering M as a point in X_n after taking a marking and rescaling the metric, this means that $M \notin W_n$.

On the other hand, the stabilizer H of M in $\operatorname{Out}(F_n)$ is isomorphic to the automorphism group of M via the homotopy equivalences between the bouquet on n circles and M. Since the quotient $M/\operatorname{Aut}(M)$ is a half-edge whose deformation space is a point, M is the unique fixed point of the group H. If there is an equivariant spine S contained in W_n , then we have $M \in S \subseteq W_n$ by Lemma 2.2, contradicting $M \notin W_n$. We conclude that W_n does not contain a spine. \Box

Remark 2.3. By taking p sufficiently large in the above proof, we see that for any $\rho > 0$, there exists some n such that the set of graphs in X_n that have at least ρn systoles does not contain an equivariant spine.

We end this section by proving the positive result that the sets V_n and V'_n are spines for X_n . The proof is the same as for Ash's well-rounded retract \mathfrak{W}_n . Recall that V_n is the set of metric graphs (of volume 1 and rank n) whose systoles are such that every component of the complement of their union is contractible (hence a finite tree without its leaves) and V'_n is the set of graphs whose systoles cover the whole graph.

Proof of Theorem 1.3. The deformation retract is performed in stages. Let U_k be the set of metric graphs such that the union of the systoles is a (possibly disconnected) graph with first Betti number at least k. Note that $U_1 = X_n$, $U_n = V_n$, and $U_k \supset U_{k+1}$ for every k. It thus suffices to construct an equivariant deformation retraction of U_k onto U_{k+1} for every k and then a deformation retraction of V_n onto V'_n .

Let $\Gamma \in U_k \setminus U_{k+1}$ for some $k \geq 1$, let S_{Γ} be the set of edges that belong to some systole and let T_{Γ} be the set of remaining edges. Let $s = s(\Gamma) < 1$ be the total length of S_{Γ} so that the total length of T_{Γ} is 1 - s > 0. For $0 \leq t \leq \log(1/s)$, we define Γ_t by rescaling the edges in S_{Γ} by a factor of e^t and those in T_{Γ} by a factor of $\frac{1-e^t s}{1-s} \geq 0$ so that the volume remains equal to 1. For t sufficiently small, the set of systoles in Γ_t stays constant because the next shortest closed geodesics in Γ are longer by a definite proportion. In particular, $\Gamma_t \in U_k \setminus U_{k+1}$ for all small enough $t \geq 0$. Let $\tau = \tau(\Gamma)$ be the supremum of times $t \in [0, \log(1/s)]$ such that $\Gamma_t \in U_k \setminus U_{k+1}$. Note that $\Gamma_{\tau} \in U_k$ since U_k is closed and Γ_t varies continuously. On the other hand, the union of the systoles in Γ_{τ} cannot be equal to a subgraph of rank exactly k otherwise we could continue the deformation for $t > \tau$, so we have $\Gamma_{\tau} \in U_{k+1}$. This is unless $\tau = \log(1/s)$, in which case the systoles in Γ_{τ} cover the whole graph and thus $\Gamma_{\tau} \in V'_n \subseteq V_n \subseteq U_{k+1}$ in that case too.

The deformation retraction $U_k \times [0,1] \to U_k$ onto U_{k+1} is defined by sending (Γ, t) to $\Gamma_{t \cdot \tau(\Gamma)}$ if $\Gamma \in U_k \setminus U_{k+1}$ and to Γ if $\Gamma \in U_{k+1}$. This map is clearly continuous, $\operatorname{Out}(F_n)$ -equivariant, equal to the identity on U_k at t = 0 and on U_{k+1} for all t, and a retract onto U_{k+1} at t = 1.

The final deformation retraction of V_n onto V'_n can be defined similarly, by shrinking all the edges that do not belong to any systole (and expanding the rest to keep the volume constant) until either the complementary components have been shrunk to points or some new systole passing through a complementary component appears. Once again, the deformation retraction is performed in stages, ordered according to the number of edges that do not belong to any systole (recall that the metric graphs in X_n are not allowed to have vertices of degree 1 or 2, so there are at most 3n - 3 edges).

3. Teichmüller space

It remains to prove Theorem 1.2 stating that the set \mathcal{W}_g of surfaces in \mathcal{T}_g whose systoles generate a finite-index subgroup in homology does not contain any equivariant spine. We simply explain how this follows from results in [FB20] and [FB23].

Proof of Theorem 1.2. Theorem 1.1 in [FB20] states that for every $\varepsilon > 0$, there exists some $g \ge 2$ and a closed hyperbolic surface X such that the systoles in X fill (so that $X \in \mathcal{V}_g$) but there are fewer than εg of them. By [FB20, Proposition 5.1], the surface constructed is such that $\operatorname{Isom}(X)$ acts transitively on a tiling of X by copies of a quadrilateral Q with three right angles and one angle of π/q for some large integer q. Taking $H = \operatorname{Isom}(X)$ as a subgroup in the extended mapping class group $\operatorname{Mod}_g^{\pm}$, we see that the locus of fixed points $\operatorname{Fix}(H)$ in \mathcal{T}_g is 1-dimensional because it is isometric to the Teichmüller space of the quotient orbifold $Q = X/\operatorname{Isom}(X)$. Trigonometric identities between the side lengths of a quadrilateral with three right angles [Bus10, p.454] imply that this space is 1-dimensional.

Now, varying the shape of Q has the effect of changing the right-angled regular 2q-gon P used to construct X into a semi-regular right-angled polygon with side lengths alternating between two values t and s(t) as in [FB23, Section 2]. If we denote the deformed surface by X_t , then the arguments in [FB23, Section 2] and [FB20, Proposition 4.1] can be easily modified to show that the systoles in X_t are either the red curves or the blue curves (or both) in the language of these papers. That is to say, the systoles in X_t are a subset of those in X. In particular, there are fewer than εg systoles in X_t for every $X_t \in Fix(H)$. Since any finite-index subgroup of $H_1(X,\mathbb{Z})$ has rank 2g, we obtain that Fix(H) is disjoint from \mathcal{W}_g as long as $\varepsilon < 2$. By Lemma 2.2, it follows that \mathcal{W}_g does not contain any Mod_g^{\pm} -equivariant spine, for otherwise $\mathcal{W}_q \cap Fix(H)$ would be non-empty. We then extend this statement to the mapping class group Mod_g . Let $H^+ \leq H$ be the index-2 subgroup of orientation-preserving isometries. Then $Q^+ = X/H^+$ is an oriented orbifold without boundary that covers Q with degree two, hence is equal to the double of Q across its boundary, i.e., a sphere with 4 cone points. The set $\operatorname{Fix}(H^+)$ is isomorphic to the Teichmüller space of Q^+ , which is isometric to the hyperbolic plane, so $\operatorname{Fix}(H^+)$ is a Teichmüller disk D containing the geodesic $L = \operatorname{Fix}(H)$.

Suppose that $S \subseteq W_g$ is a Mod_g-equivariant spine for \mathcal{T}_g . Since W_g is disjoint from L, so is S. By Lemma 2.2, $S \cap D$ is a deformation retract of D, so it is connected, hence contained in one of the two half planes bounded by L. On the other hand, $S \cap D$ is invariant under the action of the stabilizer Kof D in Mod_g. This stabilizer K contains a copy of the pure mapping class group $\operatorname{PMod}(Q^+)$ since all homeomorphisms of Q^+ fixing the cone points lift to X. In turn, $\operatorname{PMod}(Q^+)$ acts on $\mathbb{H}^2 \cong \mathcal{T}(Q^+) \cong D$ as the principal congruence subgroup of level two $\Gamma(2)$, a finite-index subgroup in $\operatorname{SL}_2(\mathbb{Z})$. In particular, K is a lattice in $\operatorname{Isom}^+(D)$, hence its limit set is all of ∂D . This contradicts the previous observation that $S \cap D$ is K-invariant and contained in a half-plane. \Box

Remark 3.1. Similarly as for Outer space, the above argument shows that for any $\rho > 0$, the set of hyperbolic surfaces that have at least ρg systoles does not contain an equivariant spine for infinitely many g.

References

- [Ash77] A. Ash, Deformation retracts with lowest possible dimension of arithmetic quotients of self-adjoint homogeneous cones, Math. Ann. **225** (1977), no. 1, 69–76.
- [Bak11] O. Baker, The Jacobian map on Outer space, PhD thesis, Cornell University, 2011.
- [Bus10] P. Buser, Geometry and spectra of compact Riemann surfaces, Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2010, Reprint of the 1992 edition.
- [CV86] M. Culler and K. Vogtmann, Moduli of graphs and automorphisms of free groups, Invent. Math. 84 (1986), no. 1, 91–119.
- [Eva79] C.W. Evans, Net structure and cages, Discrete Math. 27 (1979), no. 2, 193–204.
- [FB20] M. Fortier Bourque, Hyperbolic surfaces with sublinearly many systoles that fill, Comment. Math. Helv. 95 (2020), 515–534.
- [FB23] _____, *The dimension of Thurston's spine*, to appear in International Mathematics Research Notices, 2023+.
- [Ji14] L. Ji, Well-rounded equivariant deformation retracts of Teichmüller spaces, Enseign. Math. 60 (2014), no. 1-2, 109–129.
- [KL10] I. Kapovich and M. Lustig, Intersection form, laminations and currents on free groups, Geom. Funct. Anal. 19 (2010), no. 5, 1426–1467.
- [Mal65] A.I. Mal'cev, On the faithful representation of infinite groups by matrices, Amer. Math. Soc. Transl. Ser. 2 45 (1965), 1–18.
- [PS08a] A. Pettet and J. Souto, Minimality of the well-rounded retract, Geom. Topol. 12 (2008), no. 3, 1543–1556.
- [PS08b] _____, The spine that was no spine, Enseign. Math. (2) **54** (2008), no. 3-4, 273–285.
- [Sou75] C. Soulé, Cohomologie de SL₃(Z), C. R. Acad. Sci. Paris Sér. A-B 280 (1975), no. 5, Ai, A251–A254.

[Thu85] W.P. Thurston, A spine for Teichmüller space, preprint, 1985.

Département de mathématiques et de statistique, Université de Montréal, 2920, chemin de la Tour, Montréal (QC), H3T 1J4, Canada *Email address:* maxime.fortier.bourque@umontreal.ca